



Nyamukondiwa, C., Chidawanyika, F., Machekano, H., Mutamiswa, R., Sands, B., Mgidiswa, N., & Wall, R. (2018). Climate variability differentially impacts thermal fitness traits in three coprophagic beetle species. *PLoS ONE*, 13(6), [e0198610].
<https://doi.org/10.1371/journal.pone.0198610>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1371/journal.pone.0198610](https://doi.org/10.1371/journal.pone.0198610)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

RESEARCH ARTICLE


Climate variability differentially impacts thermal fitness traits in three coprophagic beetle species

Casper Nyamukondiwa^{1*}, Frank Chidawanyika^{2,3a}, Honest Machekano¹, Reyard Mutamiswa^{1a}, Bryony Sands⁴, Neludo Mgidiswa¹, Richard Wall⁴

1 Department of Biological Sciences and Biotechnology, Botswana International University of Science and Technology (BIUST), Palapye, Botswana, **2** Agricultural Research Council, Plant Protection Research Institute, Weeds Division, Hilton, South Africa, **3** School of Lifesciences, University of KwaZulu-Natal, Pietermaritzburg, South Africa, **4** School of Biological Sciences, University of Bristol, United Kingdom

^a Current address: Department of Zoology and Entomology, University of the Free State, Bloemfontein, South Africa

* nyamukondiwac@biust.ac.bw



Abstract

While the impacts of extreme and rising mean temperatures are well documented, increased thermal variability associated with climate change may also threaten ectotherm fitness and survival, but remains poorly explored. Using three wild collected coprophagic species *Copris elphenor*, *Metacatharsius opacus* and *Scarabaeus zambeziensis*, we explored the effects of thermal amplitude around the mean on thermal tolerance. Using standardized protocols, we measured traits of high- (critical thermal maxima [CT_{max}] and heat knockdown time [HKDT]) and -low temperature tolerance (critical thermal minima [CT_{min}], chill coma recovery time [CCRT] and supercooling points [SCPs]) following variable temperature pulses (5, 10, 15 and 20°C) around the mean (27°C). Our results show that increased temperature variability may offset basal and plastic responses to temperature and differs across species and metrics tested. Furthermore, we also show differential effects of body mass, body water content (BWC) and body lipid content (BLC) on traits of thermal tolerance. For example, body mass significantly influenced *C. elphenor* and *S. zambeziensis* CT_{max} and *S. zambeziensis* HKDT but not CT_{min} and CCRT. BWC significantly affected *M. opacus* and *C. elphenor* CT_{max} and in only *M. opacus* HKDT, CT_{min} and CCRT. Similarly, BLC only had a significant effect for *M. opacus* CT_{min}. These results suggest differential and species dependent effects of climate variability of thermal fitness traits. It is therefore likely that the ecological services provided by these species may be constrained in the face of climate change. This implies that, to develop more realistic predictions for the effects of climate change on insect biodiversity and ecosystem function, thermal variability is a significant determinant.

OPEN ACCESS

Citation: Nyamukondiwa C, Chidawanyika F, Machekano H, Mutamiswa R, Sands B, Mgidiswa N, et al. (2018) Climate variability differentially impacts thermal fitness traits in three coprophagic beetle species. PLoS ONE 13(6): e0198610. <https://doi.org/10.1371/journal.pone.0198610>

Editor: Tony Robillard, Museum National d'Histoire Naturelle, FRANCE

Received: November 9, 2017

Accepted: May 22, 2018

Published: June 6, 2018

Copyright: © 2018 Nyamukondiwa et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are available from <https://osf.io/8dg5u/> (DOI: [10.17605/OSF.IO/8DG5U](https://doi.org/10.17605/OSF.IO/8DG5U)).

Funding: Research was funded through BIUST internal research grant REF: BIUST/ds/r&i/17/2016 to CN.

Competing interests: The authors declare no competing interests and agree to this submission.

Introduction

The frequency and magnitude of extreme thermal events, including deviations from the mean environmental temperatures, is increasing due to climate change [1, 2]. These changes in both seasonal and diurnal temperature fluctuations can pose significant physiological challenges for many species [3,4], with ecological consequences that may have direct implications for biodiversity conservation [5–8]. For ectothermic arthropods, the ability to withstand both acute and chronic thermal variability is even more critical, as body temperature is largely dependent on ambient temperature [9–11]. Deviations from optimal ranges not only affect physiological processes (e.g. metabolic rates), but also key activities such as locomotion [12,13], mating, and reproductive success [14,15] and ultimately survival [16,17]. As such, the ability of organisms to operate within varying magnitudes of temperature fluctuation may be a critical trait in the face of climate change.

Phenotypic plasticity is an adaptive response associated with thermal history, that can reduce potential vulnerability to thermal extremes [8,18] both within and across generations [19,20]. For example, rapid hardening, which is a process in which an acute exposure to a sub-lethal temperature can improve subsequent performance and survival in otherwise lethal thermal conditions [21–24]. Acclimation through a chronic prior exposure to conditions similar to a future stressful environment, can improve survival and performance [25,26]. This acclimation can be realized in periods ranging from a few days to even months [11,27,28]. In many cases, such phenotypic plasticity leads only to transient physiological and biochemical responses [29–31], nevertheless, it still optimizes key life history traits during otherwise deleterious environmental conditions. Under longer timescales plasticity can lead to local genetic adaptation through changes in allele frequency [32–33].

Whilst phenotypic plasticity in response to thermal variability is pervasive in nature, several factors mediate its occurrence among different insect species. Diet quality and starvation [34–36,16], ontogeny [37,38], body size [39,40] and age [41,16] are all known to influence biochemical processes and subsequent thermal performance or survival. Indeed, strong links have been reported between body size and physiological traits such as body water and lipid content [9,42,40] with concomitant impact on climatic stress resistance [43–45,36,46]. Apart from insect physiological attributes, there is increasing evidence of how the pattern of thermal variability can also influence both constitutive (basal) and acquired (plastic) thermal responses. For example, rapid thermal variability, typical of that occurring under climate change scenarios, may exacerbate the impact of warming [47–53].

The effects of thermal history pose challenges for studies of the effects of thermal variability, because most use laboratory insect populations, typically reared under constant temperatures [54–56]. Furthermore, most studies focus on changes in mean temperatures excluding the magnitude of thermal fluctuations [57]. Thus, more robust studies on thermal variability around the mean rather than variations of the mean itself warrant investigation, in order to better understand the likely impacts of climate change on insect fitness (see [7]). As a result, here we use ecologically relevant standardised protocols (see [58,59]) to determine climate stress resistance among field-collected coprophagic beetles.

Coprophagic beetles (Coleoptera: Scarabaeidae) perform important ecosystem services such as secondary seed dispersal and nutrient cycling through degradation and burying of dung [60–62]. Furthermore, dung colonizing beetles serve as important biological control agents of intestinal parasites [63] and several pest flies [64,65] through rapid removal of the dung in the field. Due to their high diversity, abundance and important ecosystem functions, dung beetles have become a focal taxon for various ecological studies including impacts of habitat degradation [61,66–69]. The functional diversity of dung beetles has also been reported as

an important buffer for such ecosystem services in disturbed agro-ecosystems [61]. In addition, dung beetles exhibit different adult body sizes across species and losses of some ecological functions have already been attributed to poor assemblages of large sized beetles [69]. Even though the factors leading to impoverished assemblages of large sized beetles are not well understood, we hypothesize that body size differentially influences vulnerability to environmental variables, as reported in other insect taxa (e.g. [43,44,40]). How changes in mean temperature and variability affect static and plastic thermal functional traits are important for elucidating insect physiological responses [7,70]. Here we use ecologically relevant protocols [58,59] to determine climate stress resistance among field-collected coprophagic beetles. Thus we test the impact of thermal variability on overall fitness, specifically, the ability of two paracoprid (tunneller) species: *Copris elphenor* Klug, 1855, *Metacatharsius opacus* Waterhouse, 1981 and one telocoprid (ball-rolling) *Scarabaeus zambeziensis* Péringuey, 1901 to tolerate low and high temperature stress across a gradient of body mass using field collected specimens. The species to conduct the study were collected from field sites in Botswana, and they were the representatives most abundant that were active at the time of sampling.

Materials and methods

Sampling site and study animal preparation

A permit was granted from the Ministry of Environment, Wildlife and tourism Botswana [number EWT 8/36/4 XXXIII (18)] to collect experimental organisms. Dung beetles were sampled directly from the field in Khumaga Village (S20.46801; E24. 51491; 918 m.a.s.l), Central District, Botswana—a semi-arid region with Kalahari sand soils. Collections were made at the onset of the rainy season (January and February 2017), the peak period of adult emergence, to increase the likelihood of collecting beetles sharing a similar thermal history. Khumaga is characterized by abundant wildlife and domestic animals which supports high coprophagic species diversity. The area has typical average annual rainfall of 34.4 mm, most of which is recorded in January/February, and average temperatures of 23.6°C (Botswana Meteorological Services). The minimum and maximum temperatures for the two sampling months was 18.5 and 46°C respectively. Dung beetles were collected using pitfall traps baited with fresh cow dung. The traps comprised several ~8000 ml plastic buckets that were covered with a wire mesh (~5cm diameter holes) and then buried so that they were flush with the ground. The traps were equipped with a plastic rain guard that was placed 20 cm above the traps to prevent flooding. Every evening, fresh dung was placed on the wire mesh to attract foraging beetles to land and fall into the trap. A UV light bulb was also mounted on each trapping station as a light source to attract the beetles. Beetle trapping was undertaken over 12 h periods (18:00–06:00 h) and the beetles were then collected from the traps each morning. The beetles were immediately sorted according to species, based on morphological features described by Davis et al [71], and stored in insulated Styrofoam containers that had soil and fresh dung for feeding *ad libitum* during transportation to the laboratory. During the sampling period, the prevailing temperature was recorded using iButtons (model DS1923, Maxim, Sunnyvale, CA, USA). The iButtons were set within freshly deposited dung and at ground level where they were lightly covered with soil to determine potential differences in dung and ground temperature.

Thermal variability treatments

In the laboratory, insects were acclimated in climate chambers (HPP 260, Memmert GmbH + Co.KG, Germany) at $65 \pm 10\%$ relative humidity (RH) under 14L:10D at three different fluctuating thermal regimes (83, 86 and 89°C) around 27°C (Fig 1) for five days. Control beetles




Fig 1. Schematic representation of acclimation treatments to variable temperatures at different amplitudes oscillating around an ambient mean temperature of 27°C. The control (80°C) was kept at benign 27°C.

<https://doi.org/10.1371/journal.pone.0198610.g001>

were placed in a chamber at constant 27°C (80), 65 ± 10% RH and 14L:10D (Fig 1). Temperature and RH in each cage was verified using iButton data loggers.

Effects of thermal variability on high temperature plasticity

Critical thermal maxima (CT_{max}). Critical thermal maximum was measured for insects from the three treatments described above using the protocol outlined by Nyamukondiwa and Terblanche [41] and Mudavanhu et al [72]. First, ten mixed sex adult beetles were numbered and individually weighed using an electronic microbalance (RADWAG® Wagi Elektroczone, Model AS220.R2) for each of the three species and temperature treatments. This was done following 12 h starvation to allow for the clearance of gut contents and reduce the possible confounding effects on temperature tolerance [41]. Thereafter, the beetles were placed in a series of insulated double-jacketed chambers ('organ pipes') connected to a programmable water bath (Lauda Eco Gold, Lauda DR.R. Wobser GMBH and Co. KG, Germany) filled with 1:1 water: propylene glycol [9]. The beetles were then first given 10 min to equilibrate at 27°C (equivalent to the rearing benign temperature) before increasing the temperature (CT_{max}) at a rate of 0.25°C min⁻¹. This was repeated twice to give a sample size of $n = 20$ for each species per each treatment. A thermocouple (type K 36 SWG) connected to a digital thermometer (53/54IIB, Fluke Cooperation, USA) was inserted into the control chamber to monitor chamber temperatures. Individual beetle body temperature was assumed to be in equilibrium with the

chamber temperature under the experimental conditions [58,72]. The CT_{max} was defined as the temperature at which each individual insect lost coordinated muscle function, consequently losing the ability to self-right or respond to mild stimuli (like prodding with a soft thermally inert camel-hair brush) [19]. After recording, individual beetles were returned to the labeled vials fitted with gauzed lids and then placed in khaki envelopes which were placed in Memmert oven (UF 160, Memmert GmbH + Co.KG, Germany) set at 60°C for 48 h. Individual beetle weight was immediately recorded thereafter and subtracted from initial body weight to determine body water content following methods by Weldon et al. [73]. Body lipid content for *M. opacus* and *C. elphenor* was investigated and correlated with CT_{max} , following modifications from Lease & Wolf [42]. Individual adult beetles were weighed in pre-weighed 50ml Eppendorf tubes, then dried in an oven (UF160, Memmert, Germany) at 60°C for 48 h. Following drying, the flies were weighed on a RADWAG microbalance (model AS 220.R2, Poland; precision 0.001mg). Thereafter, 1.5ml of diethyl ether was added to each tube and then gently agitated at 250 rpm for 24h at 37°C using orbital shaker. The diethyl ether was then removed from the tubes, and the flies dried again at 60°C for 24h, before reweighing. The lipid content of each beetle was calculated by subtracting the lipid free dry mass from the initial beetle dry mass [42].

Heat knockdown time (HKDT). After exposure to the thermal variability treatments described in the previous section, HKDTs for the three beetle species were assayed as outlined by Weldon et al [73]. Ten beetles were weighed individually and placed in numbered 2 ml Eppendorf tubes (*M. opacus*) and 30ml polypropylene vials (*C. elphenor* and *S. zambeziensis*) and placed in a climate chamber set at $48 \pm 0.5^\circ\text{C}$ ($65 \pm 10\%$ RH) connected to a camera (HD Covert Network Camera, DS-2CD6412FWD-20, Hikvision Digital Technology Co., Ltd, China) linked to a computer. This was repeated two times to yield sample sizes of $n = 20$ for each species and temperature treatment. All observations from the climate chamber were recorded from a video recording on the computer screen. In this study, HKDT was defined as the time (in minutes) at which an individual beetle lost activity due to heat in the climate chamber as observed from the camera.

Effects of thermal variability on low temperature plasticity

Critical thermal minima (CT_{min}). For comparative low temperature tolerance of *C. elphenor*, *M. opacus*, and *S. zambeziensis* across all treatments, critical thermal minima (CT_{min}) were measured using standardized protocol as outlined by Nyamukondiwa and Terblanche [41]; Mudavanhu et al [72]. As in the CT_{max} experiments, ten replicate beetles were weighed individually and placed in numbered 'organ pipes' connected to a programmable water bath before decreasing the temperature at a rate of $0.25^\circ\text{C}/\text{min}$ until their CT_{min} were recorded. This was also repeated twice to yield sample sizes of $n = 20$ per species. In this study, CT_{min} was regarded as the temperature at which each individual insect lost coordinated muscle function, consequently losing the ability to respond to mild stimuli (e.g. gentle prodding). The BLC of *M. opacus* and *C. elphenor* was determined following previous protocol [42], and its relationship with CT_{min} was calculated.

Chill Coma Recovery Time (CCRT). Chill Coma Recovery Time experiments were assayed as outlined by Weldon et al [73]. As in the CT_{min} experiments, individual weights of *C. elphenor*, *M. opacus*, and *S. zambeziensis* were first recorded as described. After weighing, ten beetles (from each species) were individually placed in 30ml polypropylene vials with gauzed screw-cap lids and then loaded into a large zip-lock bag which was submerged into a water bath (Systronix, Scientific, South Africa) filled with 1:1 water: propylene glycol set at 0°C for 1 h. This temperature by time treatment has been previously reported to elicit chill-coma in

other insect taxa (see [73,74]). Following 1 h treatment at chill-coma temperature, the vials were immediately removed from the water bath and placed in a climate chamber set at $27 \pm 1^\circ\text{C}$, $65 \pm 10\%$ RH for recovery. The chamber was connected to a video recording camera which was linked to a computer for recording observations. This was repeated two times to yield sample sizes of $n = 20$ per treatment. CCRT was defined as the time (in mins) required for an adult to stand upright on its legs following recovery from chill-coma [75].

Supercooling points (SCPs). Supercooling points for the three beetle species were assayed following the protocol of Nyamukondiwa et al [76]. Sixteen beetles of each species were individually placed into 30 ml polypropylene vials with gauzed screw-cap lids. Each beetle was placed in contact with the tip of a type-T copper-constantan thermocouple (762–1121, Cambridge, UK), inserted through the gauzed lid of the vial and both the beetle and thermocouple were secured in contact by a cotton wool. Thermocouples were connected to one of two 8-channel Picotech TC-08 (Pico Technology, Cambridge, UK) thermocouple interfaces and temperatures were recorded at 1s intervals using PicoLog software for windows (Pico Technology, Cambridge, UK). In all treatments, experiments started by holding individual insects at 15°C for 10 mins (for insects' temperature equilibration) before decreasing the temperature at a rate of $0.5^\circ\text{C min}^{-1}$ until SCPs were recorded. SCP for each individual was determined as the lowest temperature recorded prior to a spike in temperature associated with the latent heat of crystallization [76].

Data analysis

Data was analyzed in STATISTICA 13.0 (Statsoft Inc., Tulsa, Oklahoma) and R version 3.3.0 [77]. SCPs, HKDT and CCRT data did not meet the assumptions of ANOVA and thus were analyzed using generalized linear models (GLZ) assuming a Gaussian distribution and a logit link function [27] in R3.3.0 statistical software. CTLs (CT_{\max} and CT_{\min}) met the linear model assumptions of constant variance and normal errors, so the data were analyzed using one-way factorial ANOVA in STATISTICA 13.0; Tukey-Kramer's *post-hoc* tests were used to separate statistically heterogeneous means. The relationship between traits of thermal tolerance and body mass, body water content (BWC) and BLC were examined using linear regression in STATISTICA. Differences in environmental ambient temperature and inside the dung were compared graphically using Origin 8 software (OriginLab Corporation[®], Northampton, Massachusetts, USA).

Results

Effects of thermal variability on high temperature plasticity

Thermal variability significantly affected CT_{\max} across all species tested ($F_{3,11} = 14.7$, $P < 0.001$). Higher thermal variability (86 and 89°C) was associated with a significantly lower high temperature tolerance, and this trend was more pronounced in *M. opacus* relative to *C. elphenor* and *S. zambeziensis* (Fig 2A). Similarly, there was a significant species ($F_{2,103} = 92.7$, $P < 0.001$) and species x treatment interaction effect ($F_{6,6} = 16.0$, $P < 0.001$) for CT_{\max} . Overall, *S. zambeziensis* had relatively higher CT_{\max} ($47.16 \pm 0.083^\circ\text{C}$) compared to *M. opacus* ($45.83 \pm 0.104^\circ\text{C}$) and *C. elphenor* ($45.85 \pm 0.058^\circ\text{C}$) across all treatments (Fig 2A).

Furthermore, thermal variability treatments ($\chi^2 = 118.6$, d.f = 3, $P < 0.001$) and species ($\chi^2 = 592.6.78$, d.f = 2, $P < 0.001$) also affected HKDT across all species. The interaction between species and treatment ($\chi^2 = 219.9$, d.f = 6, $P < 0.001$) also influenced HKDT. *Scara-baeus zambeziensis* had the highest HKDT indicating enhanced high temperature tolerance (Fig 2B). However, higher thermal variability had contrasting effects on the different species. While increased magnitude of temperature variability generally was associated with higher




Fig 2. Effect of temperature fluctuation (acclimation at temperatures pulsating around a benign of 27°C: 0 = 80°C (unacclimated), 3 = 83°C, 6 = 86°C and 9 = 89°C) on (A) Critical thermal maxima and (B) Heat knock down time for *M. opacus*, *C. elphenor* and *S. zambesianus*. Vertical bars denote ±95% confidence limits. Means with the same letter are not statistically different.

<https://doi.org/10.1371/journal.pone.0198610.g002>

HKDT in *C. elphenor* (51.99±1.830 minutes) and *S. zambesianus*, (75.50±4.345 minutes) it was lower in *M. opacus* (12.20±0.372 minutes) (Fig 2B).

Table 1. The relationship between traits of high and low temperature tolerance and body mass. Analysis was performed using linear regression in STATISTICA. Dung beetle species body mass were correlated with traits of temperature tolerance independently and then all species combined. *R* represents the coefficient of correlation.

Traits of high temperature	<i>r</i>	<i>p</i>
Critical Thermal Maxima (CT_{max}) vs Body mass		
All species	0.2211	0.0895
<i>M. opacus</i>	-0.3639	0.1147
<i>C. elphenor</i>	-0.4506	0.0462
<i>S. zambeianus</i>	0.4969	0.0258
Heat Knockdown Time (HKDT) vs Body mass		
All species	0.0628	0.6338
<i>M. opacus</i>	0.5598	0.0055
<i>C. elphenor</i>	0.0394	0.8689
<i>S. zambeianus</i>	-0.6497	0.0019
Traits of low temperature		
Critical Thermal Minima (CT_{min}) vs Body mass		
All species	-0.2256	0.0830
<i>M. opacus</i>	0.3419	0.1401
<i>C. elphenor</i>	-0.1307	0.5828
<i>S. zambeianus</i>	0.0840	0.7248
Chill Coma Recovery Time (CCRT) vs Body mass		
All species	0.2936	0.0280
<i>M. opacus</i>	0.2171	0.3580
<i>C. elphenor</i>	-0.2905	0.2140
<i>S. zambeianus</i>	0.0395	0.9136

<https://doi.org/10.1371/journal.pone.0198610.t001>

The relationship between traits of high temperature tolerance versus species body mass had contrasting results (Table 1). Generally, when all species were pooled together, body mass was not significantly related to high temperature tolerance. However, when species were analysed individually, body mass was significantly related to *C. elphenor* and *S. zambeianus* CT_{max} and *S. zambeianus* HKDT (Table 1). The effects of BWC on heat tolerance were more apparent both across and within species (Table 2). Due to limitations in the numbers of beetles available, it was not possible to examine the effect of BLC on all species, and this was only considered for *M. opacus* and *C. elphenor*. Nevertheless, BLC was not significantly correlated with high temperature tolerance (measured as CT_{max}) (Table 3).

Effects of thermal variability on low temperature plasticity

Thermal variability had no significant relationships with traits of low temperature tolerance including CT_{min} ($F_{3,0.65} = 1.25$, $P > 0.05$). However, it had a significant relationship with CCRT ($\chi^2 = 135.7$, d.f. = 3, $P < 0.001$) and SCPs ($\chi^2 = 13.6$, d.f. = 3, $P < 0.001$). Nevertheless, higher thermal amplitudes significantly compromised CT_{min} relative to the control (80°C), a trend most apparent in *S. zambeianus* (Fig 3A).

In contrast, higher thermal fluctuations improved CCRT in both *M. opacus* and *C. elphenor*, unlike in *S. zambeianus* (Fig 3B). Similarly, SCPs were significantly enhanced by increased thermal amplitude in both *M. opacus* and *C. elphenor*, a trend not apparent in *S. zambeianus* (Fig 3C). Furthermore, species appeared to play a significant effect on CT_{min} ($F_{2,72} = 92.9$, $P < 0.001$) and CCRT ($\chi^2 = 370.5$ d.f. = 2, $P < 0.001$) but not SCPs ($\chi^2 = 3.48$, d.f. = 2, $P > 0.05$). *Scarabaeus zambeianus* had the most compromised CT_{min} compared to the other

Table 2. The relationship between traits of high and low temperature tolerance and body water content (BWC). Analysis was performed using linear regression in STATISTICA. Dung beetle species BWC were correlated with traits of temperature tolerance independently and then all species combined. *R* represents the coefficient of correlation.

Traits of high temperature	<i>r</i>	<i>p</i>
Critical Thermal Maxima (CT _{max})		
All species	0.2593	0.00005
<i>M. opacus</i>	0.2879	0.0096
<i>C. elphenor</i>	-0.3109	0.005
<i>S. zambesianus</i>	-0.1793	0.1114
Heat Knockdown Time (HKDT)		
All species	0.4843	0.0000
<i>M. opacus</i>	0.4746	0.00001
<i>C. elphenor</i>	0.0504	0.6568
<i>S. zambesianus</i>	-0.1619	0.1515
Traits of low temperature		
Critical Thermal Minima (CT _{min})		
All species	0.0279	0.6670
<i>M. opacus</i>	0.3892	0.0004
<i>C. elphenor</i>	-0.2034	0.0703
<i>S. zambesianus</i>	-0.0445	0.6953
Chill Coma Recovery Time (CCRT)		
All species	0.5157	0.0000
<i>M. opacus</i>	0.3608	0.0010
<i>C. elphenor</i>	0.2023	0.0719
<i>S. zambesianus</i>	-0.0740	0.5143

<https://doi.org/10.1371/journal.pone.0198610.t002>

beetle species *M. opacus* and *C. elphenor*, which were not different from each other (Fig 3A). *Metacatharsius opacus* had a significantly shorter recovery time than the other beetle species following chill coma (Fig 3B).

Overall, body mass had contrasting relationships with low temperature tolerance (Table 1). Within species, body mass was not significantly related to CCRT, unlike when all species were pooled together (Table 1). Similarly, the relationship between BWC and low temperature tolerance was less pronounced, showing only significant correlation between CCRT and BWC when all species were combined, and a significant correlation for *M. opacus* for both CT_{min} and CCRT. As in high temperature versus BLC correlations, the relationship between BLC and low temperature tolerance was only undertaken for *M. opacus* and *C. elphenor*. Nevertheless,

Table 3. The relationship between body lipid content and traits of high (CT_{max}) and low (CT_{min}). Analyses were performed using linear regression in STATISTICA. Species body mass (*M. opacus* and *C. elphenor*) were correlated with traits of temperature tolerance independently and then all species combined.

Traits of temperature	<i>r</i>	<i>p</i>
Critical Thermal Maxima (CT _{max}) vs Lipid content		
All species	0.2108	0.1916
<i>M. opacus</i>	0.1022	0.6681
<i>C. elphenor</i>	-0.2264	0.3372
Critical Thermal Minima (CT _{min}) vs Body water content		
All species	0.2940	0.0656
<i>M. opacus</i>	0.6385	0.0024
<i>C. elphenor</i>	-0.4466	0.0484

<https://doi.org/10.1371/journal.pone.0198610.t003>




Fig 3. Effects of temperature fluctuations (acclimation at temperatures pulsating around a benign of 27°C: 0 = 80°C (unacclimated), 3 = 83°C, 6 = 86°C and 9 = 89°C) on (A) Critical thermal minima, (B) Chill coma recovery time and, (C) Supercooling points for *M. opacus*, *C. elphenor* and *S. zambeianus*. Vertical bars denote $\pm 95\%$ confidence limits. Means with the same letter are not statistically different.

<https://doi.org/10.1371/journal.pone.0198610.g003>

BLC had a significant relationship with low temperature tolerance (measured as CT_{min}) for both *M. opacus* and *C. elphenor* (Table 3).

Microclimate temperature recordings revealed largely similar thermal conditions within the dung and the ambient environment ($\pm 2^\circ\text{C}$) with a few exceptions where dung temperature was higher (Fig 4).

Discussion

Here the effects of exposure to thermal variability are reported for natural field-derived populations of coprophagic beetles. Our results show that higher amplitudes generally constrain the thermal fitness traits tested, with a few exceptions, highlighting the complexity of insect responses to temperature means and variability. Notably, CT_{max} and HKDT results generally showed that increased variability in temperature may compromise high temperature tolerance. This suggests that, under projected increase in thermal variability, the coprophagic species studied here may be at more risk of high temperature mortality, a trend that was apparent in *M. opacus* and *C. elphenor* (for CT_{max}) and *S. zambeianus* for HKDT (Fig 2A and 2B). These results are consistent with findings by Terblanche et al [7] who demonstrated loss of basal and plastic responses to temperature in response to increased amplitude of temperature fluctuations.

Given that global climate change predictions suggest an increased frequency and variability of stressful temperatures [78,79], understanding the effects of these changes to insect population dynamics is important [80]. Related studies have looked at the effects of thermal amplitude on growth [81], fertility [82], density dependent net reproductive rate [57], development rate [83], stress resistance [84] and vector borne disease transmission [85]. However, none has considered natural populations of coprophagic species, with a view of understanding effects of climate variability on their activity and provision of ecosystem services. In addition, more recent studies have focused on how magnitudes of the variation impact species thermal performance curves [47,49, reviewed in 80]. Our current findings are therefore the first to show that increased amplitudes of thermal variations, typical of those predicted under global climate change, may have significant fitness costs to coprophagic beetles. This is likely to affect their provision of important ecosystem services [86,65].

Some studies suggest temperature variability may independently affect fitness traits, owing to the non-linearity of thermal reaction norms or thermal performance curves (see [56,70]). Our results also show that, while thermal amplitude plays a significant role in fitness, species also differ significantly in their response to temperature. Among the three species, *S. zambeianus* appeared more tolerant to high temperature basally, compared to the other two beetle species. This further supports the idea that arthropod responses to climate variability are species-specific, and the rate of anthropogenic climate change may exceed that which may be compensated through genetic and evolutionary plastic mechanisms among the three different beetle species (e.g. [7]). The findings also suggest that *M. opacus* may be at more risk of high temperature mortality following higher thermal fluctuations compared to *C. elphenor* and *S. zambeianus*. Previous studies have also documented that while plasticity to high temperature tolerance is an evident and near-ubiquitous feature of insect thermal physiology [9,87], the benefits may not be adequate to provide complete compensation for global climate change




Fig 4. Differences in microclimate temperature records between ambient and 'inside-dung' during a one-week dung pat decomposition period in Khumaga village. Temperature was recorded using i-Buttons (DS1923, Maxim, Sunnyvale, CA, USA) at 15 minutes sampling interval for 5 days (January 19–23 2017). One i-Button was immersed in the middle of a freshly dropped cow dung pat, while the other one was placed in a shaded environment, ~1 m above the ground.

<https://doi.org/10.1371/journal.pone.0198610.g004>

[88,87]. This therefore suggests that other mechanisms for compensation, e.g. behavioural modification and evolution may offer more benefit. However, it seems likely that many species may not be able to use behaviour e.g. tracking preferred microenvironments in space under prolonged environmental stress (e.g. [89]). This notion appears true for the dung beetle species tested here since ambient and dung temperature records were similar (Fig 4). It has been suggested that telocoprid dung beetle species (e.g. *S. zambezianus* studied here) withstand high temperatures by frequently mounting the dung to avoid excessive ground heat (of up to 60°C) during the day [90]. However, since our ground temperature recordings closely matched that of the dung it is unlikely that this ball-rolling behaviour would protect the species that were tested here from excessive heat. Furthermore, our study included two paracoprid species *C. elphenor* and *M. opacus*. The microclimate climate data recorded here suggest that physiological mechanisms to mitigate excessive heat are more important for their survival and preservation of key activities. However, the species tested here are nocturnal and perhaps gain significant performance and survival advantage due to cooler temperatures at night than day. It is possible that this nocturnal foraging behavior might be an evolutionary response to evade excessive day temperatures.

The correlation between body mass and high temperature tolerance was trait and species dependant. For example, in *S. zambezianus*, CT_{max} was positively correlated with body mass whilst HKDT was negative. The current results therefore contradict previous study by Nyamukondiwa and Terblanche [41] which found no evidence for the body mass constraints on thermal tolerance. However, our study supports studies where body mass was shown to correlate with heat tolerance [40,45,91]. We also found that increased heat tolerance was associated with increased BWC, as previously reported [45], perhaps due to enhanced resistance to desiccation [92]. This may be because insect mortality at high temperatures is closely associated with rapid water loss leading to desiccation [9,92]. Thus, insects with higher BWC may take more time to

desiccate, and may have improved survival at high temperature through the link between the two stressors (see [93]). We did not find any correlation between traits of high temperature and BLC for the beetle species tested here, indicating that body lipid content may not play a role in high temperature tolerance.

Thermal variability had more dramatic effects on CCRT compared to CT_{min} and SCPs. The results therefore highlight differences in sensitivity to thermal variability among related traits of thermal tolerance. For *S. zambeziensis* CT_{min} , higher thermal amplitudes also compromised low temperature tolerance (Fig 3A). It is also interesting to note that *S. zambeziensis* had the highest lower activity temperature limit (CT_{min}), which perhaps relates to its superior high temperature tolerance compared to the other species (see Fig 2A and 2B), since this may have come at a cost of fitness at low temperature (CT_{min}) [94]. Indeed, similar fitness costs have been reported in other insect taxa, [95–97]. Similar modest thermal fluctuations have been observed to affect many insect life history traits including mating behaviour and fertility (see [9]). Similarly, Marshall and Sinclair [98] showed that at stressful low temperatures, *D. melanogaster* trades off immediate survival for future reproductive output. Thus, thermal variability reported here may also have significant life history trade-offs for the beetle species tested. Coupled with individual reduced fitness, thermal variability under changing climates may therefore have far reaching population level consequences [47]. Nevertheless, exact trade-offs resulting from the magnitude of the thermal variability in *M. opacus*, *C. elphenor* and *S. zambeziensis* warrants future investigation.

Previous studies have indicated that ectotherms are already experiencing sub-lethal low temperatures, e.g. chill coma and CT_{min} in natural environments [47], and consistently compromise fitness traits under increased magnitude of temperature changes [9,70]. While the direct and indirect effects of increased thermal variability have been appreciated by evolutionary ecologists [56,70], mechanistic models for surviving variable stressful environments have been limited. These mechanisms however remain significant in explaining how organisms likely cope with imminent and increased thermal variability associated with climate change. Without compensatory physiological or behavioural modifications to fluctuating thermal regimes, current findings thus suggest fitness losses for the beetle species tested here. Measurement of traits of low temperature *vis* CT_{min} , CCRT and chill coma also reveal significant patterns of species local low temperature adaptation and compensatory responses under thermal variability (see [9]). However, Donat and Alexander [99] suggest cold events may be less likely as opposed to extreme high temperature events under climate change. Thus survival of species under variable high temperature may be more imminent as opposed to low temperatures. Nevertheless, while low temperature variability may not be that critical here, it has shifted key life history traits, seasonal timing, and population phenologies of many insect species [100].

Body mass generally had little impact on low temperature tolerance traits (Table 1) even when traits for all species were pooled together, in particular for CCRT. This result corroborates with Nyamukondiwa and Terblanche [41] who showed no effects of body mass on insect CTLs to activity. Similarly, the relationship between high temperature tolerance and BWC did not follow a specific pattern. For example, BWC appeared to be positively correlated with CT_{min} and CCRT for *M. opacus*, and when species were pooled together for CCRT. This suggests taxa and trait dependent effects of varying thermal amplitudes to low temperature tolerance. Our results also showed significant correlations between traits of low temperature and BLC. Indeed, [101] elaborated on the role of metabolites, including carbohydrates (i.e. trehalose), free amino acids, lipids, osmoprotectants and polyols are upregulated during low temperature conditions. Our results therefore suggest lipid accumulation may be an essential mechanism sustaining low temperature survival by insects faced with increased magnitude of low temperature fluctuations.

Age and body condition e.g. nutrition are among some of the factors which may affect physiological and biochemical processes in insects, including traits of thermal tolerance measured here [9,34,38]. Thus, one confounding effect in our study here is that, age and nutritional history of the test organisms was not strictly controlled for. Nevertheless, our assessments of body mass, BWC and BLC and their correlation with thermal tolerance traits may be indicative of body condition which can correlate with age. In conclusion our results show that coprophagic insects *M. opacus*, *C. elphenor* and *S. zambeziensis* may suffer fitness losses (for both high and low temperature traits) in response to increased temperature variability with climate change. Tolerance levels varied significantly across variability (8) levels, and likely depend on the trait in question and the species. Field studies have reported that pesticide residues in the dung of treated animals may have deleterious consequences on different developmental stages of dung beetles [102,103]. Thus loss of fitness with thermal variability reported here, coupled with reports on the effects of pesticides (e.g. [103]), represent a significant burden to conservation of dung beetle ecological services.

Acknowledgments

We appreciate logistical support from the Ministry of Environment, Wildlife and Tourism Botswana and Michael Flyman for the research permits, Botswana International University of Science and Technology (BIUST), and University of KwaZulu-Natal. We are also grateful to the Insect Physiological Ecology research group (BIUST) for assistance in running experiments and several anonymous reviewers for their comments on an earlier version of this manuscript.

Author Contributions

Conceptualization: Casper Nyamukondiwa.

Data curation: Honest Machekano, Reyard Mutamiswa, Bryony Sands, Neludo Mgidiswa.

Formal analysis: Casper Nyamukondiwa, Frank Chidawanyika, Honest Machekano, Reyard Mutamiswa.

Funding acquisition: Casper Nyamukondiwa.

Investigation: Honest Machekano, Reyard Mutamiswa, Neludo Mgidiswa.

Methodology: Bryony Sands.

Project administration: Casper Nyamukondiwa.

Resources: Casper Nyamukondiwa, Richard Wall.

Software: Casper Nyamukondiwa.

Supervision: Casper Nyamukondiwa, Richard Wall.

Validation: Casper Nyamukondiwa, Frank Chidawanyika, Bryony Sands.

Visualization: Honest Machekano.

Writing – original draft: Casper Nyamukondiwa.

Writing – review & editing: Casper Nyamukondiwa, Frank Chidawanyika, Honest Machekano, Reyard Mutamiswa, Bryony Sands, Neludo Mgidiswa, Richard Wall.

References

1. Diffenbaugh NS, Pal JS, Trapp RJ, Giorgi F (2005) Fine-scale processes regulate the response of extreme events to global climate change. *Proceedings of the National Academy of Sciences of the*

- United States of America 102: 15774–15778. <https://doi.org/10.1073/pnas.0506042102> PMID: 16236722
2. Diffenbaugh NS, Field CB (2013) Changes in ecologically critical terrestrial climate conditions. *Science* 341: 486–492. <https://doi.org/10.1126/science.1237123> PMID: 23908225
3. Helmuth B (2009) From cells to coastlines: how can we use physiology to forecast the impacts of climate change? *Journal of Experimental Biology* 212: 753–760. <https://doi.org/10.1242/jeb.023861> PMID: 19251989
4. Sunday JM, Bates AE, Kearney MR, Colwel RK, Dulvy NK, Longino JT et al. (2014) Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. *Proceedings of the National Academy of Sciences of the United States of America* 111: 5610–5615. <https://doi.org/10.1073/pnas.1316145111> PMID: 24616528
5. Sinclair BJ, Addo-Bediako A, Chown SL (2003) Climatic variability and the evolution of insect freeze tolerance. *Biological reviews of the Cambridge Philosophical Society* 78: 181–195. PMID: 12803420
6. Chown SL, Terblanche JS (2006) Physiological Diversity in Insects: Ecological and Evolutionary Contexts. *Advances in Insect Physiology* 33: 50–152. [https://doi.org/10.1016/S0065-2806\(06\)33002-0](https://doi.org/10.1016/S0065-2806(06)33002-0) PMID: 19212462
7. Terblanche JS, Nyamukondiwa C, Kleynhans E (2010) Thermal variability alters climatic stress resistance and plastic responses in a globally invasive pest, the Mediterranean fruit fly (*Ceratitidis capitata*). *Entomologia Experimentalis et Applicata* 137: 304–315.
8. Gunderson AR, Stillman JH (2015) Plasticity in thermal tolerance has limited potential to buffer ectotherms from global warming. *Proceedings of the Royal Society of London B* 282: 1808. <https://doi.org/10.1098/rspb.2015.0401> PMID: 25994676
9. Chown SL, Nicolson SW (2004) *Insect physiological ecology: mechanisms and patterns*. Oxford University Press. New York.
10. Speight MR, Hunter MD, Watt AD (2008) *Ecology of Insects: Concepts and Applications*. Wiley-Blackwell, West Sussex, UK Chichester, U.K. 626 pp.
11. Fischer K, Dierks A, Franke K, Geister TL, Liska M, Winter S et al. (2010) Environmental effects on temperature stress resistance in the tropical butterfly *Bicyclus anynana*. *PLOS ONE* 5: e15284. <https://doi.org/10.1371/journal.pone.0015284> PMID: 21187968
12. Clusella-Trullas S, Terblanche JS, Chown SL (2010) Phenotypic Plasticity of locomotion performance in the seed harvester *Messor capensis* (Formicidae) *Physiological and Biochemical Zoology* 83: 519–530. <https://doi.org/10.1086/651387> PMID: 20367320
13. Lachenicht MW, Clusella-Trullas S, Boardman L, Le Roux C, Terblanche JS (2010) Effects of acclimation temperature on thermal tolerance, locomotion performance and respiratory metabolism in *Acheta domesticus* L. (Orthoptera: Gryllidae). *Journal of Insect Physiology* 56: 822–830. <https://doi.org/10.1016/j.jinsphys.2010.02.010> PMID: 20197070
14. Piyaphongkul J, Pritchard J, Bale J (2012) Heat stress impedes development and lowers fecundity of the brown planthopper *Nilaparvata lugens* (Stål). *PLoS ONE* 7: e47413. <https://doi.org/10.1371/journal.pone.0047413> PMID: 23071803
15. Dick CA, Rank NE, McCarthy M, McWeeney S, Hollis D, Dahlhoff E P (2013) Effects of temperature variation on male behavior and mating success in a Montane beetle. *Physiological and Biochemical Zoology* 86: 432–440. <https://doi.org/10.1086/671462> PMID: 23799837
16. Chidawanyika F, Nyamukondiwa C, Strathie L, Fischer K (2017) Effects of thermal regimes, starvation and age on heat tolerance of the Parthenon beetle *Zygogramma bicolorata* (Coleoptera: Chrysomelidae) following dynamic and static protocols. *PLoS One* 12(1): e0169371. <https://doi.org/10.1371/journal.pone.0169371> PMID: 28052099
17. Mutamiswa R, Chidawanyika F, Nyamukondiwa C (2017) Comparative assessment of the thermal tolerance of spotted stemborer, *Chilo Partellus* (Lepidoptera: Crambidae) and its larval parasitoid, *Cotesia Sesamiae* (Hymenoptera: Braconidae). *Insect Science*. 00: 1–14.
18. Valladares F, Matesanz S, Guilhaumon F, Araujo MB, Balaguer L, Benito-Garzon M et al. (2014) The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecology Letters* 17: 1351–1364. <https://doi.org/10.1111/ele.12348> PMID: 25205436
19. Nyamukondiwa C, Terblanche JS (2010) Within-generation variation of critical thermal limits in adult Mediterranean and Natal fruit flies *Ceratitidis capitata* and *Ceratitidis rosa*: thermal history affects short-term responses to temperature. *Physiological Entomology* 35: 255–264.
20. Zhou Z-S, Rasmann S, Li M, Guo J-Y, Chen H-S, Wan F-H (2013) Cold temperatures increase cold hardiness in the next generation *Ophraella communa* beetles. *PLoS ONE* 8: e74760. <https://doi.org/10.1371/journal.pone.0074760> PMID: 24098666

21. Sejerkilde M, Jesper G, Sørensen JG, Loeschcke V (2003) Effects of cold- and heat hardening on thermal resistance in *Drosophila melanogaster*. *Journal of Insect Physiology* 49: 719–726. PMID: [12880651](#)
22. Kellett M, Hoffmann AA, McKechnie SW (2005) Hardening capacity in the *Drosophila melanogaster* species group is constrained by basal thermotolerance. *Functional Ecology* 19: 853–858.
23. Stotter RL, Terblanche JS (2009) Low temperature tolerance of false codling moth *Thaumotobia leucotreta* (Meyrick) (Lepidoptera: Tortricidae) in South Africa. *Journal of Thermal Biology* 34: 320–325
24. Chidawanyika F, Terblanche JS (2011a) Rapid thermal responses and thermal tolerance in adult codling moth *Cydia pomonella* (Lepidoptera: Tortricidae). *Journal of Insect Physiology* 57: 108–117. <https://doi.org/10.1016/j.jinsphys.2010.09.013> PMID: [20933517](#)
25. Huey RB, Berrigan D (1996) Testing evolutionary hypotheses of acclimation. In Johnston IA and Bennett AE (eds.), *Phenotypic and evolutionary adaptation to temperature*, pp. 205–237. U. Cambridge Press, Cambridge, U.K.
26. Geister TL, Fischer K (2007) Testing the beneficial acclimation hypothesis: temperature effects on mating success in a butterfly. *Behavioural Ecology* 18: 658–664.
27. Chidawanyika F, Terblanche JS (2011b) Costs and benefits of thermal acclimation for codling moth, *Cydia pomonella* (Lepidoptera: Tortricidae): implications for pest control and the sterile insect release programme. *Evolutionary Applications* 4: 534–544. <https://doi.org/10.1111/j.1752-4571.2010.00168.x> PMID: [25568003](#)
28. Kristensen TN, Hoffmann AA, Overgaard J, Sørensen JG, Hallas R, Loeschcke V (2008) Costs and benefits of cold acclimation in field-released *Drosophila*. *Proceedings of the National Academy of Sciences of the United States of America* 105: 216–221. <https://doi.org/10.1073/pnas.0708074105> PMID: [18162547](#)
29. Boardman L, Sørensen JG, Terblanche JS (2013) Physiological responses to fluctuating thermal and hydration regimes in the chill susceptible insect, *Thaumotobia leucotreta*. *Journal of Insect Physiology* 59: 781–794. <https://doi.org/10.1016/j.jinsphys.2013.05.005> PMID: [23684741](#)
30. Boardman L, Sørensen JG, Terblanche JS (2015) Physiological and molecular mechanisms associated with cross tolerance between hypoxia and low temperature in *Thaumotobia leucotreta*. *Journal of Insect Physiology* 82: 75–84. <https://doi.org/10.1016/j.jinsphys.2015.09.001> PMID: [26376454](#)
31. Sgrò CM, Terblanche JS, Hoffmann AA (2016) What Can Plasticity Contribute to Insect Responses to Climate Change? *Annual review of Entomology* 61: 433–451. <https://doi.org/10.1146/annurev-ento-010715-023859> PMID: [26667379](#)
32. Angilletta MJ Jr Bennett AF, Guderley H Navas CA, Seebacher F Wilson RS (2006) Coadaptation: A Unifying Principle in Evolutionary Thermal Biology. *Physiological and Biochemical Zoology* 79: 282–294. <https://doi.org/10.1086/499990> PMID: [16555188](#)
33. Gerken AR, Eller OC, Hahn DA, Morgan TJ (2015) Constraints, independence, and evolution of thermal plasticity: Probing genetic architecture of long- and short-term thermal acclimation. *Proceedings of the National Academy of Sciences of the United States of America* 112: 4399–4404. <https://doi.org/10.1073/pnas.1503456112> PMID: [25805817](#)
34. Shreve SM, Yi SX, Lee RE (2007) Increased dietary cholesterol enhances cold tolerance in *Drosophila melanogaster*. *Cryoletters* 28: 33–37. PMID: [17369960](#)
35. Diamond SE, Kingsolver JG (2010) Environmental dependence of thermal reaction norms: host plant quality can reverse the temperature-size rule. *American Naturalist* 175: 1–10. <https://doi.org/10.1086/648602> PMID: [19911984](#)
36. Scharf I, Wexler Y, MacMillan HA, Presman S, Simson E, Rosenstein S (2016) The negative effect of starvation and the positive effect of mild thermal stress on thermal tolerance of the red flour beetle, *Tribolium castaneum*. *Science of Nature* 103: 20. <https://doi.org/10.1007/s00114-016-1344-5> PMID: [26888763](#)
37. Marais K, Terblanche JS, Chown SL (2009) Life stage-related differences in hardening and acclimation of thermal tolerance traits in the kelp fly, *Paractora dreuxi* (Diptera, Helcomyzidae). *Journal of Insect Physiology* 55: 336–43. <https://doi.org/10.1016/j.jinsphys.2008.11.016> PMID: [19171152](#)
38. Bowler K, Terblanche JS (2008) Insect thermal tolerance: what is the role of ontogeny, ageing and senescence? *Biological Reviews* 83: 339–355. PMID: [18979595](#)
39. Spicer JI, Gaston KJ (1999) *Physiological diversity and its ecological implications*. Oxford, UK: Blackwell Science.
40. Klockmann M, Gunter F, Fischer K (2017) Heat resistance throughout ontogeny: body size constrains thermal tolerance. *Global Change Biology* 23: 686–696. <https://doi.org/10.1111/gcb.13407> PMID: [27371939](#)

41. Nyamukondiwa C, Terblanche JS (2009) Thermal tolerance in adult Mediterranean and Natal fruit flies (*Ceratitis capitata* and *Ceratitis rosa*): effects of age, gender and feeding status. *Journal of Thermal Biology* 34: 406–414.
42. Lease HM, Wolf BO (2011) Lipid content of terrestrial arthropods in relation to body size, phylogeny, ontogeny and sex. *Physiological Entomology* 36: 29–38.
43. Le Lagadec M, Chown S, Scholtz C (1998) Desiccation resistance and water balance in southern African keratin beetles (Coleoptera, Trogidae): the influence of body size and habitat. *Journal of Comparative Physiology B* 168: 112–122.
44. Chown SL, Le Lagadec MD, Scholtz CH (1999) Partitioning variance in a physiological trait: desiccation resistance in keratin beetles (Coleoptera, Trogidae). *Functional Ecology* 13: 838–844.
45. Oberg EW, Toro ID, Pelini SL (2012) Characterization of the thermal tolerances of forest ants of New England. *Insectes Sociaux* 59: 167–174.
46. Mitchell KA, Boardman L, Clusella-Trullas S, Terblanche JS (2017) Effects of nutrient and water restriction on thermal tolerance: A test of mechanisms and hypotheses. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 212: 15–23. <https://doi.org/10.1016/j.cbpa.2017.06.019> PMID: 28676379
47. Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC et al. (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Science USA* 105: 6668–6672.
48. Paaïjms KP, Heinig RL, Seliga RA, Blanford JI, Blanford S, Murdock CC et al. (2013) Temperature variation makes ectotherms more sensitive to climate change. *Global Change Biology* 19: 2373–2380. <https://doi.org/10.1111/gcb.12240> PMID: 23630036
49. Kingsolver JG, Diamond SE, Buckley LB, (2013) Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology* 27: 1415–1423.
50. Estay SA, Lima M, Bozinovic F (2014) The role of temperature variability on insect performance and population dynamics in a warming world. *Oikos* 123: 131–140.
51. Dowd WW, King FA, Denny MW (2015) Thermal variation, thermal extremes and the physiological performance of individuals. *Journal of Experimental Biology* 218: 1956–1967. <https://doi.org/10.1242/jeb.114926> PMID: 26085672
52. Ma FZ, Lu ZC, Wang R, Wan FH (2014) Heritability and evolutionary potential in thermal tolerance traits in the invasive Mediterranean cryptic species of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *PLoS ONE* 9:e103279. <https://doi.org/10.1371/journal.pone.0103279> PMID: 25054554
53. Shah AA, Gill BA, Encalada AC, Flecker AS, Funk WC, Guayasamin JM et al. (2017) Climate variability predicts thermal limits of aquatic insects across elevation and latitude. *Functional Ecology* 31: 2118–2127.
54. Scheiner SM (2002) Selection experiments and the study of phenotypic plasticity. *Journal of Evolutionary Biology* 15: 889–898.
55. Schulte PM, Healy TM, Fanguie NA (2011) Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology* 51: 691–702. <https://doi.org/10.1093/icb/ucr097> PMID: 21841184
56. Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS et al. (2014) Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society of London B* 281:20132612.
57. Clavijo-Baquet S, Boher F, Ziegler L, Martel SI, Estay SA, Bozinovic F (2014) Differential responses to thermal variation between fitness metrics. *Scientific Reports* 4: 5349. <https://doi.org/10.1038/srep05349> PMID: 24954717
58. Terblanche JS, Deere JA, Clusella-Trullas S, Janion C, Chown SL (2007) Critical thermal limits depend on methodological context. *Proceedings of the Royal Society of London Series B, Biological Sciences* 274: 2935–2943.
59. Terblanche JS, Hoffmann AA, Mitchell KA, Rako L, le Roux PC, Chown SL (2011) Ecologically relevant measures of tolerance to potentially lethal temperatures. *Journal of Experimental Biology* 214: 3713–3725. <https://doi.org/10.1242/jeb.061283> PMID: 22031735
60. Yamada D, Imura O, Shi K, Shibuya T (2007) Effect of tunneler dung beetles on cattle dung decomposition, soil nutrients and herbage growth. *Grassland Science* 53: 121–129.
61. Beynon SA, Mann DJ, Slade EM, Lewis OT (2012) Species-rich dung beetle communities buffer ecosystem services in perturbed agro-ecosystems. *Journal of Applied Ecology* 49: 1365–1372.
62. Beynon SA, Wainwright WA, Christie M (2015) The application of an ecosystem services framework to estimate the economic value of dung beetles to the U.K. cattle industry. *Ecological Entomology* 40: 124–135.

63. Sands B, Wall R (2016) Dung beetles reduce livestock gastrointestinal parasite availability on pasture. *Journal of Applied Ecology* 54: 1180–1189.
64. Bergstrom RC, Maki LR, Werner BA (1976). Small dung beetles as biological control agents: laboratory studies of beetle action on trichostrongylid eggs in sheep and cattle feces. *Proceedings of the Helminthological Society of Washington* 43: 171–174.
65. Nichols E, Spector S, Louzada J, Larsen T, Amezcua S, Favila ME (2008) Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* 141: 1461–1474.
66. Nichols E, Larsen T, Spector S, Davis AL, Escobar F, Favila M et al. (2007) Dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. *Biological Conservation* 137: 1–19.
67. Jacobs CT, Scholtz CH, Escobar F, Davis ALV (2010) How might intensification of farming influence dung beetle diversity (Coleoptera: Scarabaeidae) in Maputo Special Reserve (Mozambique)? *Journal of Insect Conservation* 14: 389–399.
68. Almeida S, Louzada J, Sperber C, Barlow J (2011) Subtle land-use change and tropical biodiversity: dung beetle communities in cerrado grasslands and exotic pastures. *Biotropica* 43: 704–710.
69. Braga RF, Korasaki V, Andresen E, Louzada J (2013) Dung beetle community and functions along a habitat-disturbance gradient in the Amazon: a rapid assessment of ecological functions associated to biodiversity. *PLoS One* 8:e57786. <https://doi.org/10.1371/journal.pone.0057786> PMID: 23460906
70. Bozinovic F, Medina NR, Alruiz JM, Cavieres G, Sabat P (2016) Thermal tolerance and survival responses to scenarios of experimental climatic change: changing thermal variability reduces the heat and cold tolerance in a fly. *Journal of Comparative Physiology* 186: 581–587. <https://doi.org/10.1007/s00360-016-0980-6> PMID: 27003422
71. Davis ALV, Frolov AV, Scholtz CH (2008) *The African Dung Beetle Genera*. Protea Book House. Pretoria, South Africa.
72. Mudavanhu P, Addison P, Conlong DE (2014) Impact of mass rearing and gamma irradiation on thermal tolerance of *Eldana saccharina*. *Entomologia Experimentalis et Applicata* 153: 55–63.
73. Weldon CW, Terblanche JS, Chown SL (2011) Time-course for attainment and reversal of acclimation to constant temperature in two *Ceratitis* species. *Journal of Thermal Biology* 36: 479–485.
74. Sinclair BJ, Coello Alvarado LE, Ferguson LV (2015) An invitation to measure insect cold tolerance: Methods, approaches, and workflow. *Journal of Thermal Biology* 53: 180–197. <https://doi.org/10.1016/j.jtherbio.2015.11.003> PMID: 26590471
75. Milton CC, Partridge L (2008) Brief carbon dioxide exposure blocks heat hardening but not cold acclimation in *Drosophila melanogaster*. *Journal of Insect Physiology* 54: 32–40. <https://doi.org/10.1016/j.jinsphys.2007.08.001> PMID: 17884085
76. Nyamukondiwa C, Weldon CW, Chown SL, le Roux PC, Terblanche JS (2013) Thermal biology, population fluctuations and implications of temperature extremes for the management of two globally significant insect pests. *Journal of Insect Physiology* 59: 1199–1211. <https://doi.org/10.1016/j.jinsphys.2013.09.004> PMID: 24080125
77. R Development Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
78. Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO (2000) Climate extremes: observations, modeling, and impacts. *Science* 289: 2068–2074. PMID: 11000103
79. Meehl GA, Tebaldi C (2004) More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305: 994–997. <https://doi.org/10.1126/science.1098704> PMID: 15310900
80. Thompson RM, Beardall J, Beringer J, Grace M, Sardina P (2013) Means and extremes: building variability into community-level climate change experiments. *Ecology Letters* 16: 799–806. <https://doi.org/10.1111/ele.12095> PMID: 23438320
81. Estay SA, Clavijo-Baquet S, Lima M, Bozinovic F (2011) Beyond average: an experimental test of temperature variability on the population dynamics of *Tribolium confusum*. *Population Ecology* 53: 53–58.
82. Ragland GJ, Kingsolver JG (2008) The effect of fluctuating temperatures on ectotherm life history traits: comparisons among geographic populations of *Wyeomyia smithii*. *Evolutionary Ecology Research* 10: 29.
83. Saunders LM, Tompkins DM, Hudson PJ (2002) Stochasticity accelerates nematode egg development. *Journal of Parasitology* 88: 1271–1272. [https://doi.org/10.1645/0022-3395\(2002\)088\[1271:SANED\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2002)088[1271:SANED]2.0.CO;2) PMID: 12537129
84. Folguera G, Bastián DA, Caers J, Rojas JM, Piulachs M-D, Belle's X et al. (2011) An experimental test of the role of environmental temperature variability on ectotherm molecular, physiological and life-

- p>history traits: implications for global warming.
- Comparative Biochemistry and Physiology A*
- 159: 242–246.
85. Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB (2010) Influence of climate on malaria transmission depends on daily temperature variation. *Proceedings of the National Academy of Science USA* 107: 15135–15139.
86. Losey JE, Vaughan M (2006) The economic value of ecological services provided by insects. *Bioscience* 56: 311–323.
87. Gunderson AR, Dillon ME, Stillman JH (2017) Estimating the benefits of plasticity in ectotherm heat tolerance under natural thermal variability. *Functional Ecology* 31: 529–1539.
88. Mitchell KA, Hoffmann AA (2017) Thermal ramping rate influences evolutionary potential and species differences for upper thermal limits in *Drosophila*. *Functional Ecology* 24: 694–700.
89. Chevin LM, Lande R, Mace GM (2010) Adaptation, plasticity and extinction in a changing environment. Towards a predictive theory. *PLoS Biology* 8: e1000357. <https://doi.org/10.1371/journal.pbio.1000357> PMID: 20463950
90. Smolka J, Baird E, Bryne MJ, Jundi B, Warrant EJ, Dacke M (2012) Dung beetles use their dung ball ball as mobile thermal refuge. *Current Biology* 22: 863–864.
91. Brans KI, Jansen M, Vanoverbeke J, Tuzun N, Stoks R, De Meester L (2017) The heat is on: Genetic adaptation to urbanization mediated by thermal tolerance and body size. *Global Change Biology*. <https://doi.org/10.1111/gcb.13784> PMID: 28614592
92. Weldon CW, Boardman L, Marlin D, Terblanche JS (2016) Physiological mechanisms of dehydration tolerance contribute to the invasion potential of *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) relative to its less widely distributed congeners. *Frontiers in Zoology* 13: 15. <https://doi.org/10.1186/s12983-016-0147-z> PMID: 27034703
93. Bubliy OA, Kristensen TN, Kellermann V, Loeschcke V (2012) Plastic responses to four environmental stresses and cross resistance in a laboratory population of *Drosophila melanogaster*. *Functional Ecology* 26: 245–253.
94. Nyamukondiwa C, Terblanche JS, Marshall KE, Sinclair BJ (2011) Basal cold but not heat tolerance constraints plasticity among *Drosophila* species (Diptera: Drosophilidae). *Journal of Evolutionary Biology* 24: 1927–1938. <https://doi.org/10.1111/j.1420-9101.2011.02324.x> PMID: 21658189
95. Hoffmann AA, Hewa-Kapuge S (2000) Acclimation for heat resistance in *Trichogramma* nr. *brassicae*: can it occur without costs? *Functional Ecology* 14: 55–60.
96. Coulson SC, Bale JS. (1992) Effect of rapid cold hardening on reproduction and survival of offspring in the housefly *Musca domestica*. *Journal of Insect Physiology* 38: 421–424.
97. Basson CH, Nyamukondiwa C, Terblanche JS (2011). Fitness costs of rapid cold-hardening in *Ceratitis capitata*. *Evolution* 66: 296–304. <https://doi.org/10.1111/j.1558-5646.2011.01419.x> PMID: 22220884
98. Marshall KE, Sinclair BJ (2010) Repeated stress exposure results in a survival-reproduction trade-off in *Drosophila melanogaster*. *Proceedings of the Royal Society of London B* 277: 963–969.
99. Donat MG, Alexander LV (2012) The shifting probability distribution of global daytime and night-time temperatures. *Geophysics Research Letters* 39: <https://doi.org/10.1029/2012GL052459>
100. Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution and Systematics* 37: 637–669.
101. Sinclair BJ (2015) Linking energetics and overwintering in temperate insects. *Journal of Thermal Biology* 54: 5–11. <https://doi.org/10.1016/j.jtherbio.2014.07.007> PMID: 26615721
102. Kruger K, Lukhele OM, Scholtz CH (1999) Survival and reproduction of *Euoniticellus intermedius* (Coleoptera: Scarabaeidae) in dung following application of cypermethrin and flumethrin pour-ons to cattle. *Bulletin of Entomological Research* 89: 543–548.
103. Mann CM, Barnes S, Offer B, Wall R (2015) Lethal and sub-lethal effects of faecal deltamethrin residues on dung-feeding insects. *Medical and Veterinary Entomology* 29: 189–195. <https://doi.org/10.1111/mve.12104> PMID: 25594879